

Exclusive double diffractive events: menu for LHC

Vladimir Alexeevich Petrov and Roman Anatolievich Rytin

Institute for High Energy Physics

142 281, Protvino, Russia

E-mail: petrov@mail.ihep.ru, ryutin@th1.ihep.su

ABSTRACT: Exclusive double diffractive events (EDDE) are considered in the framework of the Regge-eikonal approach and perturbative calculations for “hard” subprocesses. Total and differential cross-sections for processes $p + p \rightarrow p + X + p$ are calculated.

KEYWORDS: Higgs Physics, Heavy Quarks Physics, Jets, Phenomenological Models.

Contents

1. Introduction	1
2. Calculations	2
3. Results for resonance production	6
4. Standard model Higgs boson production	7
5. Heavy quarkonium production	8
6. Radion production	8
7. EDD dijet production	10
8. Conclusions	11

1. Introduction

LHC collaborations aimed at working in low and high p_T regimes related to typical undulatory (diffractive) and corpuscular (point-like) behaviours of the corresponding cross-sections may offer a very exciting possibility to observe an interplay of both regimes [1]. In theory the “hard part” can be (hopefully) treated with perturbative methods whilst the “soft” one is definitely nonperturbative.

Below we give several examples of such an interplay: exclusive particle production by diffractively scattered protons, i.e. the processes $p + p \rightarrow p + X + p$, where $+$ means also a rapidity gap and X represents a particle or a system of particles consisting of or strongly coupled to the two-gluon state.

This process is related to the dominant amplitude of exclusive two-gluon production. Driving mechanism of the diffractive processes is the Pomeron. Data on the total cross-sections demands unambiguously for the Pomeron with larger-than-one intercept, thereof the need in “unitarisation”.

As will be seen below, EDDE gives us unique experimental possibilities for particle searches and investigations of diffraction itself. This is due to several advantages of the process: a) clear signature of the process; b) possibility to use “missing mass method”, that improve the mass resolution; c) background is strongly suppressed; d) spin-parity analysis of the central system can be done; e) interesting measurements concerning the interplay between “soft” and “hard” scales are possible.




Figure 1: The “bare” amplitude of the process $p + p \rightarrow p + X + p$.




Figure 2: Unitarization of the process $p + p \rightarrow p + X + p$.

2. Calculations

In figures 1 and 2 we illustrate in detail the process $p + p \rightarrow p + X + p$. Off-shell proton-gluon amplitudes in figure 1 are treated by the method developed in ref. [2], which is based on the extension of Regge-eikonal approach, and successfully used for the description of the HERA data [3].

The amplitude of the process $p + \bar{p} \rightarrow p + X + p$ can be obtained in the following way (see figures 1 and 2). The first step is to calculate the “bare” amplitude T_X , which is depicted in figure 1. The “hard” part is the usual gluon-gluon fusion process calculated by perturbative methods in the Standard Model or its extensions. “Soft” amplitudes $T_{1,2}$ are obtained in the Regge-eikonal approach. The second step is the unitarization procedure, that takes into account initial and final state interactions (see figure 2).

We use the following kinematics, which corresponds to the double Regge limit. It is convenient to use light-cone components $(+, -, \perp)$. The components of momenta of the hadrons in figure 1 are

$$\begin{aligned}
 p_1 &= \left(\sqrt{\frac{s}{2}}, \frac{m^2}{\sqrt{2s}}, \mathbf{0} \right) \\
 p_2 &= \left(\frac{m^2}{\sqrt{2s}}, \sqrt{\frac{s}{2}}, \mathbf{0} \right) \\
 p'_1 &= \left((1 - \xi_1) \sqrt{\frac{s}{2}}, \frac{\Delta_1^2 + m^2}{(1 - \xi_1) \sqrt{2s}}, -\Delta_1 \right) \\
 p'_2 &= \left(\frac{\Delta_2^2 + m^2}{(1 - \xi_2) \sqrt{2s}}, (1 - \xi_2) \sqrt{\frac{s}{2}}, -\Delta_2 \right) \\
 q &= (q_+, q_-, \mathbf{q}), \\
 q_1 &= q + p_1 - p'_1 = q + \Delta_1, \\
 q_2 &= -q + p_2 - p'_2 = -q + \Delta_2,
 \end{aligned} \tag{2.1}$$

$\xi_{1,2}$ are fractions of protons momenta carried by gluons. For two-dimensional transverse vectors we use boldface type. From the above notations we can obtain the relations:

$$\begin{aligned}
 t_{1,2} = \Delta_{1,2}^2 &\simeq -\frac{\Delta_{1,2}^2(1 + \xi_{1,2}) + \xi_{1,2}^2 m^2}{1 - \xi_{1,2}} \simeq -\Delta_{1,2}^2, \quad \xi_{1,2} \rightarrow 0 \\
 \cos \phi_0 &= \frac{\Delta_1 \Delta_2}{|\Delta_1| |\Delta_2|} \\
 M_X^2 &\simeq \xi_1 \xi_2 s + t_1 + t_2 - 2\sqrt{t_1 t_2} \cos \phi_0 \\
 (p_1 + q)^2 &\simeq m^2 + q^2 + \sqrt{2s} q_- = s_1 \\
 (p_2 - q)^2 &\simeq m^2 + q^2 - \sqrt{2s} q_+ = s_2.
 \end{aligned} \tag{2.2}$$

Physical region of diffractive events with two rapidity gaps is defined by the following kinematical cuts:

$$0.01 \text{ GeV}^2 \leq |t_{1,2}| \leq 1 \text{ GeV}^2, \tag{2.3}$$

$$\xi_{\min} \simeq \frac{M_X^2}{s \xi_{\max}} \leq \xi_{1,2} \leq \xi_{\max} = 0.1, \tag{2.4}$$

$$\begin{aligned}
 (\sqrt{-t_1} - \sqrt{-t_2})^2 &\leq \kappa \leq (\sqrt{-t_1} + \sqrt{-t_2})^2 \\
 \kappa &= \xi_1 \xi_2 s - M_X^2 \ll M_X^2
 \end{aligned} \tag{2.5}$$

The discussion on the choice of cuts (2.3)–(2.5) for diffractive events and references to other authors were given in [4, 5]. We can write the relations in terms of $y_{1,2}$ and y_X (rapidities of hadrons and the system X correspondingly). For instance:

$$\begin{aligned}
 \xi_{1,2} &\simeq \frac{M_X}{\sqrt{s}} e^{\pm y_X}, \\
 |y_X| &\leq y_0 = \ln \left(\frac{\sqrt{s} \xi_{\max}}{M_X} \right), \\
 y_0 &\simeq 2.5 \quad \text{for } \sqrt{s} = 14 \text{ TeV}, \quad M_X = O(100 \text{ GeV}), \\
 |y_{1,2}| &= \frac{1}{2} \ln \frac{(1 - \xi_{1,2})^2 s}{m^2 - t_{1,2}} \geq 9
 \end{aligned} \tag{2.6}$$

In standard terms the amplitude corresponds to the so-called nonfactorized scheme [4]. The contribution of the diagram depicted in figure 2 is obtained by integrating over all internal loop momenta. It was shown in [4], that the leading contribution arises from the region of the integration, where momentum q is ‘‘Glauber-like’’, i.e. of the order $(\mathbf{k}_+ m^2 / \sqrt{s}, \mathbf{k}_- m^2 / \sqrt{s}, \mathbf{k}m)$, where \mathbf{k} ’s are of the order 1. The detailed consideration of the loop integral like

$$\int \frac{d^4 q}{(2\pi)^4} \frac{f(q, p_1, p_2, \Delta_1, \Delta_2)}{(q^2 + i0)(q_1^2 + i0)(q_2^2 + i0)} \tag{2.7}$$

shows that the main contribution comes from the poles at

$$\begin{aligned}
 q_1^2 &= \sqrt{2s} \xi_1 q_- - \mathbf{q}_1^2 = 0, \\
 q_2^2 &= -\sqrt{2s} \xi_2 q_+ - \mathbf{q}_2^2 = 0.
 \end{aligned}$$

In this case

$$q = \left(-\frac{\mathbf{q}_2^2}{\xi_2 \sqrt{2s}}, \frac{\mathbf{q}_1^2}{\xi_1 \sqrt{2s}}, \mathbf{q} \right), \quad (2.8)$$

where

$$\begin{aligned} \mathbf{q}_1^2 &= \mathbf{q}^2 + \Delta_1^2 + 2|\mathbf{q}||\Delta_1| \cos \left(\phi + \frac{\phi_0}{2} \right), \\ \mathbf{q}_2^2 &= \mathbf{q}^2 + \Delta_2^2 - 2|\mathbf{q}||\Delta_2| \cos \left(\phi - \frac{\phi_0}{2} \right) \end{aligned}$$

Taking the general form for T -amplitudes that satisfy identities

$$q^\alpha T_{\mu\alpha}^D = 0, \quad q_i^\mu T_{\mu\alpha}^D = 0, \quad (2.9)$$

and neglecting terms of the order $o(\xi_i)$, the following expression is found at $|t_i| \leq 1 \text{ GeV}^2$:

$$\begin{aligned} T_{\mu\alpha}^D(p, q, q_i) &= \left(G_{\mu\alpha} - \frac{P_\mu^{q_i} P_\alpha^q}{P^{q_i} P^q} \right) T_{gp \rightarrow gp}^D(s_i, t_i, qq_i) \\ G_{\mu\alpha} &= g_{\mu\alpha} - \frac{q_{i,\mu} q_\alpha}{qq_i}, \\ P_\mu^{q_i} &= p_\mu - \frac{pq_i}{qq_i} q_\mu, \\ P_\alpha^q &= p_\alpha - \frac{pq}{qq_i} q_{i,\alpha}. \end{aligned} \quad (2.10)$$

For $T_{gp \rightarrow gp}^D$ we use the Regge-eikonal approach [2, 6]. At small t_i it takes the form of the Born approximation, i.e. Regge factor:

$$T_{gp \rightarrow gp}^D(s_i, t_i, qq_i) = c_{gp} \left(e^{-i\frac{\pi}{2}} \frac{s_i - qq_i - m^2}{s_0 - qq_i - m^2} \right)^{\alpha_P(t_i)} e^{b_0 t_i}, \quad b_0 = \frac{1}{4} \left(\frac{r_{pp}^2}{2} + r_{gp}^2 \right), \quad (2.11)$$

where

$$\begin{aligned} \alpha_P(0) &= 1.203, \quad \alpha'_P(0) = 0.094 \text{ GeV}^{-2}, \\ r_{pp}^2 &= 2.477 \text{ GeV}^{-2} \end{aligned} \quad (2.12)$$

are fixed parameters for the ‘‘hard’’ Pomeron [6], which have been obtained from the global fit to the data on diffractive $pp(p\bar{p})$ scattering. Parameters

$$c_{gp} \simeq 3.5, \quad r_{gp}^2 = 2.54 \text{ GeV}^{-2} \quad (2.13)$$

are defined from fitting the HERA data on elastic J/Ψ production [7], which can be found in [8] and will be published elsewhere. It is shown that the Born term corresponding to the ‘‘hard’’ Pomeron with parameters (2.12) of the trajectory gives the main contribution to the amplitude. The upper bound for the constant $c_{gp}^{up} \simeq 2.3(3.3)$ can be also estimated from the exclusive double diffractive di-jet production at Tevatron (see (7.4)), if we take CDF cuts and the upper limit for the exclusive total di-jet cross-section [9]. The effective value $c_{gp} = 2.3$ corresponds to the case, when the Sudakov suppression factor is absorbed into the constant, and $c_{gp} = 3.3$ is obtained when taking into account this factor explicitly.

The full “bare” amplitude looks as follows:

$$T_{pp \rightarrow pXp} \simeq \int \frac{d^4q}{(2\pi)^4} \frac{8F^{\mu\nu}(q_1, q_2) T_{\mu\alpha}^D(p_1, q, q_1) T_{\nu\alpha}^D(p_2, -q, q_2)}{(q^2 + i0)(q_1^2 + i0)(q_2^2 + i0)}, \quad (2.14)$$

where

$$F^{\mu\nu}(q_1, q_2) = \left(g^{\mu\nu} - \frac{q_2^\mu q_1^\nu}{M_X^2} \right) F_{gg \rightarrow X}.$$

Factor 8 arises from the colour index contraction. Let $l^2 = -q^2 \simeq \mathbf{q}^2$, $y_X = \langle y_X \rangle = 0$ and contract all the tensor indices, then the integral (2.14) takes the form

$$T_{pp \rightarrow pXp} \simeq c_{gp}^2 e^{b(t_1+t_2)} \frac{\pi}{(2\pi)^2} \left(-\frac{s}{M_X^2} \right)^{\alpha_P(0)} \cdot 8F_{gg \rightarrow X} \cdot I, \quad (2.15)$$

$$b = \alpha'_P(0) \ln \left(\frac{\sqrt{s}}{M_X} \right) + b_0, \quad (2.16)$$

$$I \simeq \int_0^{M_X^2} \frac{dl^2}{l^4} \left(\frac{l^2}{s_0 - m^2 + l^2/2} \right)^{2\alpha_P(0)}, \quad (2.17)$$

where $s_0 - m^2 \simeq 1 \text{ GeV}^2$ is the scale parameter of the model that is used in the global fitting of the data on $pp(p\bar{p})$ scattering for on-shell amplitudes [6]. It remains fixed in the present calculations. It worth nothing that the “rescattering corrections” for the off-shell gluon-proton amplitudes, T^D , are small (in accordance with a general analysis in ref. [2]). Contrary to this, the “outer” corrections (see eq. (2.20)) are significant.

If we take into account the emission of virtual “soft” gluons, while prohibiting the real ones, that could fill rapidity gaps, it results in the Sudakov-like suppression [10]:

$$F_s(l^2) = \exp \left[-\frac{3}{2\pi} \int_{l^2}^{M_X^2/4} \frac{dp_T^2}{p_T^2} \alpha_s(p_T^2) \ln \left(\frac{M_X^2}{4p_T^2} \right) \right], \quad (2.18)$$

and in the new value of the integral (2.17):

$$I_s \simeq \int_0^{M_X^2} \frac{dl^2}{l^4} F_s(l^2) \left(\frac{l^2}{s_0 - m^2 + l^2/2} \right)^{2\alpha_P(0)}. \quad (2.19)$$

In this case the total cross-section becomes smaller, than without the factor F_s . It plays significant role for large M_X .

Unitarity corrections can be estimated from the elastic pp scattering by the method depicted in figure 2, where

$$\begin{aligned} T_X &= T_{pp \rightarrow pXp}, \\ V(s, \mathbf{q}_T) &= 4s(2\pi)^2 \delta^2(\mathbf{q}_T) + 4s \int d^2\mathbf{b} e^{i\mathbf{q}_T \cdot \mathbf{b}} \left[e^{i\delta_{pp \rightarrow pp}} - 1 \right], \\ T_X^{\text{Unit.}}(p_1, p_2, \Delta_1, \Delta_2) &= \frac{1}{16ss'} \int \frac{d^2\mathbf{q}_T}{(2\pi)^2} \frac{d^2\mathbf{q}'_T}{(2\pi)^2} V(s, \mathbf{q}_T) \times \\ &\quad \times T_X(p_1 - q_T, p_2 + q_T, \Delta_{1T}, \Delta_{2T}) \cdots V(s', \mathbf{q}'_T), \\ \Delta_{1T} &= \Delta_1 - q_T - q'_T, \\ \Delta_{2T} &= \Delta_2 + q_T + q'_T, \end{aligned} \quad (2.20)$$

and $\delta_{pp \rightarrow pp}$ can be found in ref. [6]. Left and right parts V represent “soft” rescattering effects for initial and final states, i.e. multi-Pomeron exchanges. These “outer” unitarity corrections reduce the integrated cross-section by the factor about 14 for the given kinematical cuts and lead to the changes in the ϕ_0 -dependence, which will be discussed in further publications.

3. Results for resonance production

We have the following expression for the differential cross-section in case of one particle production:

$$\frac{d\sigma}{dt_1 dt_2 d\xi_1 d\xi_2} = \frac{\pi |T_{pp \rightarrow pXp}^{Unit.}|^2}{8s(2\pi)^5 \sqrt{-\lambda}} \quad \lambda = \kappa^2 + 2(t_1 + t_2)\kappa + (t_1 - t_2)^2 \leq 0. \quad (3.1)$$

By partial integrating (3.1) we obtain t and ξ distributions. The first result of our calculations is depicted in the figure 3. The antishrinkage of the diffraction peak with increasing mass M_X is the direct consequence of the existence of the additional hard scale M_X , which makes the interaction radius smaller. The ξ distribution is shown in figure 4.

We can use the following replacement to obtain the cross-section for the EDD process $p + p \rightarrow p + X + p$:

$$|F_{gg \rightarrow X}|^2 \rightarrow 4\pi M_X \Gamma(X \rightarrow gg). \quad (3.2)$$

It is possible to simplify calculations after reduction of (3.1) to

$$\frac{d\sigma}{dt_1 dt_2 d\phi_0} \simeq \frac{\pi |T_{pp \rightarrow pXp}^{Unit.}|_{y_X=0}^2}{8s^2(2\pi)^5} \Delta y_X, \quad (3.3)$$

where $\Delta y_X = 2y_0$, ϕ_0 is the azimuthal angle between outgoing protons.




Figure 3: t -distribution $d\sigma/dt/\sigma_{tot}$ of the process $p + p \rightarrow p + X + p$ for masses of the system X equal to 20 GeV (solid one) and 50 GeV (dashed one).




Figure 4: ξ -distribution $d\sigma/d\xi/\sigma_{tot}$ of the process $p + p \rightarrow p + X + p$ for $M_X = 100$ GeV.

4. Standard model Higgs boson production

For the Standard Model Higgs boson [11]

$$F_{gg \rightarrow H}^0 = M_H^2 \frac{\alpha_s}{2\pi} \sqrt{\frac{G_F}{\sqrt{2}}} f(\eta), \tag{4.1}$$

$$f(\eta) = \frac{1}{\eta} \left\{ 1 + \frac{1}{2} \left(1 - \frac{1}{\eta} \right) \left[Li_2 \left(\frac{2}{1 - \sqrt{1 - \frac{1}{\eta}} - i0} \right) + Li_2 \left(\frac{2}{1 + \sqrt{1 - \frac{1}{\eta}} + i0} \right) \right] \right\},$$

$$|F_{gg \rightarrow H}|^2 \rightarrow 1.5 |F_{gg \rightarrow H}^0|^2, \tag{4.2}$$

where $\eta = M_H^2/4m_t^2$, G_F is the Fermi constant, m_t is the top quark mass. NLO K-factor 1.5 for the $gg \rightarrow H$ process is included to the final answer.

Numerical results are the following [12]

c_{gp}	M_H (GeV)	$\sigma_{p+p \rightarrow p+H+p}$ (fb)			
		LHC		TeVatron	
		no Sud. suppr.	Sud. suppr.	no Sud. suppr.	Sud. suppr.
3.5	100 \rightarrow 500	110 \rightarrow 57	4.6 \rightarrow 0.14	12 \rightarrow 0.4	0.5 \rightarrow 0.001
2.3	100 \rightarrow 500	20 \rightarrow 11	–	2.2 \rightarrow 0.08	–
3.3	100 \rightarrow 500	–	3.6 \rightarrow 0.11	–	0.4 \rightarrow 0.0009

We consider four different cases only for Higgs boson as illustration of the Sudakov suppressing action. In other examples we take $c_{gp} = 3.3$ with Sudakov-like suppression according to the CDF data estimations to obtain the lower value of cross-sections at LHC and Tevatron. For this case our result is quite close to the one of [10], where the value of the total cross-section is about 3fb. In both cases the most important suppressing in the mass region $M_H > 100$ GeV is due to (perturbative) Sudakov factors, while the nonperturbative (absorbitive) factors play relatively minor role.

Results of other authors were considered in details in [13]. Here we refer to the highest cross-section $0.25 \rightarrow 0.4$ pb for $M_H = 100$ GeV at LHC energies that was obtained in ref. [5]. A nonfactorized form of the amplitude and a “QCD inspired” model for $gp \rightarrow gp$ amplitudes were used, taking into account the nonperturbative proton wave functions. Even if we multiply the result of ref. [5] by the suppressing factor, it will be larger than ours. This could serve as the indication of the role of nonperturbative effects. Our model is based on the Regge-eikonal approach for the amplitudes, which is primordially nonperturbative, normalized to the data from HERA on $\gamma p \rightarrow J/\Psi p$ [7] and improved by the CDF data on the exclusive di-jet production [9].

To estimate the signal to QCD background ratio for $b\bar{b}$ signal we use the standard expression for $gg \rightarrow b\bar{b}$ amplitude and assumptions [14]-[17]:

- possibility to separate final $b\bar{b}$ quark jets from gluon jets. If we cannot do it, it will increase the background by two orders of magnitude. The efficiency of b-tagging is supposed to be 50%.

- suppression due to the absence of colour-octet $b\bar{b}$ final states
- suppression of light fermion pair production, when $J_{z,tot} = 0$ (see also [18, 19])
- cut $E_T > 50 \text{ GeV}$ ($\theta \geq 60^\circ$), since the cross-section of EDD $b\bar{b}$ jet production strongly decreases with E_T (see formulae (7.2)).

The theoretical result of our numerical estimations is

$$\frac{\text{Signal}(pp \rightarrow pHp \rightarrow pb\bar{b}p)}{\text{QCD background}} \geq 3.8 \frac{\text{GeV}}{\Delta M}, \quad (4.3)$$

where ΔM is the mass resolution of the detector and $M_H \simeq 115 \text{ GeV}$, which can reach $0.01M_H$ due to application of the “missing mass method”. Similar result was strictly obtained in [14, 15]. Under the above circumstances the total efficiency at the integrated luminosity 30 fb^{-1} is $\sim 10\%$ and numerically estimated significance of the event is about 3σ , which is close to the one in $\gamma\gamma$ decay mode.

5. Heavy quarkonium production

Results for $\chi_{c0,b0}$ EDD production were obtained recently by some authors [14, 16, 20] in different approaches. To obtain the total cross-sections in the model considered in the present paper let us substitute widths of these states into (3.2).

$$\Gamma(\chi_{b0} \rightarrow gg) \simeq \Gamma_0(\chi_{b0} \rightarrow gg) \left(1 + 9.8 \frac{\alpha_S}{\pi}\right) = 550 \text{ keV} \quad (\text{see [14, 16] for details}), \quad (5.1)$$

where the width is set to the lattice result $\Gamma_0(\chi_{b0} \rightarrow gg) = 354 \text{ keV}$ [21]. After replacement we obtain for LHC and TeVatron:

$$\sigma_{pp \rightarrow p+\chi_{b0}+p} \simeq 1.3 \text{ nb}, \quad \sqrt{s} = 14 \text{ TeV}, \quad ((2.3), (2.4) \text{ cuts}), \quad (5.2)$$

$$\sigma_{pp \rightarrow p+\chi_{b0}+p} \simeq 160 \text{ pb}, \quad \sqrt{s} = 1.8 \text{ TeV}, \quad (\text{CDF cuts}). \quad (5.3)$$

The same procedure can be done for χ_{c0} . Taking the total width $\Gamma(\chi_{c0} \rightarrow gg) \simeq 14.9 \text{ MeV}$ [22] we obtain

$$\sigma_{pp \rightarrow p+\chi_{c0}+p} \simeq 4 \mu\text{b}, \quad \sqrt{s} = 14 \text{ TeV}, \quad ((2.3), (2.4) \text{ cuts}), \quad (5.4)$$

$$\sigma_{pp \rightarrow p+\chi_{c0}+p} \simeq 600 \text{ nb}, \quad \sqrt{s} = 1.8 \text{ TeV}, \quad (\text{CDF cuts}). \quad (5.5)$$

6. Radion production

Now there is a great interest to the multidimensional properties of the space-time. One of the models was proposed by Randall and Sundrum [23]. We have considered the case of one compact extra-dimension. In this case we have additional scalar particle Radion, that reflects the existence of an extra dimension and represents the field of the “distance” oscillations between the branes along the extra dimension. Since Radion has the same quantum numbers as the Higgs boson, they can mix [24]. After mixing we have two




Figure 5: The total cross-section (in fb) of the process $p + p \rightarrow p + h^* + p$ versus Higgs boson(h^*) mass for $c_{gp} = 3.3$ with Sudakov-like suppression at LHC. Parameters of RS1 model are shown.

mass eigenstates, which could be observed experimentally. For the EDDE the following replacements in (4.1) should be done:

$$f(\eta) \rightarrow a_{34}f(\eta_{h^*}) + 7\gamma b \quad \text{for } h^*, \tag{6.1}$$

$$f(\eta) \rightarrow \gamma(a_{12}f(\eta_{r^*}) + 7a) \quad \text{for } r^*, \tag{6.2}$$

where $\eta_{h^*,r^*} = m_{h^*,r^*}^2/4m_t^2$ and other parameters are obtained from formulae in the appendix A of [24]:

$$\gamma = \frac{v}{\Lambda_\phi}, \quad v = 246\text{GeV is the Higgs VEV, } \Lambda_\phi \text{ is the radion VEV,} \tag{6.3}$$

$$Z^2 = 1 - 6\xi\gamma^2(1 + 6\xi), \quad \tan 2\theta = 12\xi\gamma Z \frac{1}{Z^2 - 36\xi^2\gamma^2 - m_r^2/m_h^2}, \tag{6.4}$$

$$a = \cos\theta/Z; \quad b = -\sin\theta/Z; \quad c = \sin\theta - 6\xi\gamma/Z; \quad d = \cos\theta + 6\xi\gamma/Z \sin\theta, \tag{6.5}$$

$$a_{12} = a + c/\gamma; \quad a_{34} = d + b\gamma, \tag{6.6}$$

$$m_{r^*}^2 = c^2m_h^2 + a^2m_r^2; \quad m_{h^*}^2 = d^2m_h^2 + b^2m_r^2, \tag{6.7}$$

$$r = ar^* + bh^*; \quad h = cr^* + dh^*. \tag{6.8}$$

Results for the total cross-sections in the case of $c_{gp} = 3.3$ with Sudakov-like suppression are depicted in figures 5 and 6 for several values of mixing parameter ξ and for the vacuum expectation value of the radion field $\Lambda_\phi = 1 \text{ TeV}$.




Figure 6: The total cross-section (in fb) of the process $p + p \rightarrow p + r^* + p$ versus Radion(r^*) mass for $c_{gp} = 3.3$ with Sudakov-like suppression at LHC. Parameters of RS1 model are shown.

7. EDD dijet production

For the EDD production of a dijet system of mass M_X in the leading order (see, for example [4]) we have:

$$|F_{gg \rightarrow X}|^2 \rightarrow \frac{144\pi^2 \alpha_S^2 M_X^4}{E_T^4}, \quad X = gg, \quad (7.1)$$

$$|F_{gg \rightarrow X}|^2 \rightarrow \frac{32\pi^2 \alpha_S^2 M_X^2 m_Q^2}{3E_T^4} \beta^2, \quad X = Q\bar{Q}, \quad \beta = \sqrt{1 - \frac{4m_Q^2}{M_X^2}}, \quad (7.2)$$

and the cross-section becomes

$$\frac{d\sigma}{dt_1 dt_2 dy_X d\kappa' dE_T^2} \simeq \frac{|T_{pp \rightarrow pjjp}^{Unit.}|^2}{2^{13} \pi^5 s^2 \kappa' \sqrt{1 - \kappa'}}, \quad (7.3)$$

where $\kappa' = 4E_T^2/M_X^2$, and $T_{pp \rightarrow pjjp}^{Unit.}$ is calculated as in the section 2 with substitutions (7.1), (7.2).

Note that all the results of this article are given for the value of the constant $c_{gp} = 3.3$, which is obtained from the upper bounds for the exclusive di-jet production at TeVatron energies [9]. Cross-sections and numerical estimations for c_{gp} at different transverse energy

cuts are the following:

$$\begin{aligned}
 E_T > 7 \text{ GeV}, & \quad \sigma < 3.7 \text{ nb}, & \quad c_{gp} < 3.3 \\
 E_T > 10 \text{ GeV}, & \quad \sigma < 0.97 \pm 0.065 \text{ (stat.)} \pm 0.272 \text{ (sys.) nb}, & \quad c_{gp} < 3.4 \\
 E_T > 25 \text{ GeV}, & \quad \sigma < 34 \pm 5 \text{ (stat.)} \pm 10 \text{ (sys.) pb}, & \quad c_{gp} < 4.2.
 \end{aligned} \tag{7.4}$$

The lowest value is close to the result, obtained by fitting the HERA data on elastic J/Ψ production. It can serve as the indication of model applicability.

From the analogous calculations for LHC with cuts (2.3), (2.4) we have:

$$\begin{aligned}
 E_T > 10 \text{ GeV}, & \quad \sigma(pp \rightarrow p + \text{jet} + \text{jet} + p) \simeq 7 \text{ nb} \\
 E_T > 25 \text{ GeV}, & \quad \sigma(pp \rightarrow p + \text{jet} + \text{jet} + p) \simeq 150 \text{ pb} \\
 E_T > 50 \text{ GeV}, & \quad \sigma(pp \rightarrow p + \text{jet} + \text{jet} + p) \simeq 8 \text{ pb}.
 \end{aligned} \tag{7.5}$$

8. Conclusions

We see from the results that there is a real possibility to use advantages of the the EDDE for investigations at LHC. Accuracy of the mass measurements could be improved by applying the missing mass method [25].

The low value of the exclusive Higgs boson production cross-section obtained in this paper is mainly due to the Sudakov suppression factor (2.18), the full validity of which is not obvious, because the confinement effects can strongly modify the “real gluon emission”. It is interesting that in spite of different models and quite different ways of account of absorbtive effects in our paper and in ref. [10], the final results appeared to be quite close.

Certainly, the cross-sections may be several times larger due to still not very well known non-perturbative factors.

In the case of heavy quarkonium and dijet EDD production cross-sections are much larger than for Higgs boson production, and some other important investigations like measurements of the azimuthal angle dependence and the diffractive pattern of the interaction could be done.

Acknowledgments

We are grateful to A. De Roeck, A. Prokudin, A. Rostovtsev, N.E. Tyurin, A. Sobol, S. Slabospitsky and participants of BLOIS2003 workshop and several CMS meetings for helpful discussions. We are also indebted to N.V. Krasnikov and V.A. Matveev, who draw our attention to the radion phenomenology. This work is supported by the Russian Foundation for Basic Research, grant no. 04-02-17299.

References

- [1] V.A. Petrov, *CMS+TOTEM: joint measurement of “soft-hard” diffractive events*, in *Proc. of the 2nd Int. Symp. LHC: Physics and Detectors*, A.N. Sissakian and Y.A. Kultchitsky eds., Dubna June 2000;
 V.A. Petrov, A.V. Prokudin, S.M. Troshin and N.E. Tyurin, *Novel features of diffraction at the LHC*, *J. Phys.* **G 27** (2001) 2225.

- [2] V.A. Petrov, *High-energy implications of extended unitarity*, in *Proceedings of the VIIth Blois Wokshop*, M. Haguenaue et al. eds., Editions Frontières, Paris 1995.
- [3] V.A. Petrov and A.V. Prokudin, *Is a soft pomeron appropriate for describing hera data?*, *Phys. Atom. Nucl.* **62** (1999) 1562
- [4] A. Berera and J.C. Collins, *Double pomeron jet cross sections*, *Nucl. Phys.* **B 474** (1996) 183 [[hep-ph/9509258](#)].
- [5] J.-R. Cudell and O.F. Hernandez, *Particle production in a hadron collider rapidity gap: the Higgs case*, *Nucl. Phys.* **B 471** (1996) 471 [[hep-ph/9511252](#)].
- [6] V.A. Petrov and A.V. Prokudin, *The first three pomerons*, *Eur. Phys. J.* **C 23** (2002) 135 [[hep-ph/0105209](#)].
- [7] H1 collaboration, S. Aid et al., *Elastic and inelastic photoproduction of J/ψ mesons at HERA*, *Nucl. Phys.* **B 472** (1996) 3 [[hep-ex/9603005](#)]; *Elastic photoproduction of J/ψ and upsilon mesons at HERA*, *Phys. Lett.* **B 483** (2000) 23 [[hep-ex/0003020](#)]; ZEUS collaboration, J. Breitweg et al., *Measurement of elastic J/ψ photoproduction at HERA*, *Z. Physik* **C 75** (1997) 215 [[hep-ex/9704013](#)]; *Measurement of elastic upsilon photoproduction at HERA*, *Phys. Lett.* **B 437** (1998) 432 [[hep-ex/9807020](#)].
- [8] V.A. Petrov, A.V. Prokudin and R.A. Ryutin, *From the exclusive photoproduction of heavy quarkonia at HERA to the EDDE at TeVatron and LHC*, [hep-ph/0404116](#).
- [9] ON BEHALF OF THE CDF collaboration, M. Gallinaro, *Diffraction physics results at CDF*, *Acta Phys. Polon.* **B35** (2004) 465 [[hep-ph/0311192](#)].
- [10] V.A. Khoze, A.D. Martin and M.G. Ryskin, *Can the Higgs be seen in rapidity gap events at the Tevatron or the LHC?*, *Eur. Phys. J.* **C 14** (2000) 525 [[hep-ph/0002072](#)]; *High p_T Higgs signal for the LHC*, *Eur. Phys. J.* **C 21** (2001) 99 [[hep-ph/0104230](#)]; V.A. Khoze, *Rapidity gap Higgs signal at the Tevatron and the LHC*, [hep-ph/0105224](#).
- [11] B.A. Kniehl, *Higgs phenomenology at one loop in the standard model*, *Phys. Rept.* **240** (1994) 211.
- [12] V.A. Petrov and R.A. Ryutin, *Exclusive double diffractive Higgs boson production at LHC*, *Eur. Phys. J.* **C 36** (2004) 509 [[hep-ph/0311024](#)].
- [13] V.A. Khoze, A.D. Martin and M.G. Ryskin, *Diffractive Higgs production: myths and reality*, *Eur. Phys. J.* **C 26** (2002) 229 [[hep-ph/0207313](#)].
- [14] V.A. Khoze, A.D. Martin and M.G. Ryskin, *Double-diffractive processes in high-resolution missing- mass experiments at the Tevatron*, *Eur. Phys. J.* **C 19** (2001) 477 [[hep-ph/0011393](#)].
- [15] A. De Roeck, V.A. Khoze, A.D. Martin, R. Orava and M.G. Ryskin, *Ways to detect a light Higgs boson at the LHC*, *Eur. Phys. J.* **C 25** (2002) 391 [[hep-ph/0207042](#)].
- [16] V.A. Khoze, A.D. Martin and M.G. Ryskin, *Prospects for new physics observations in diffractive processes at the LHC and Tevatron*, *Eur. Phys. J.* **C 23** (2002) 311 [[hep-ph/0111078](#)].
- [17] A.D. Martin, M.G. Ryskin and V.A. Khoze, *Dijet hadroproduction with rapidity gaps and QCD double logarithmic effects*, *Phys. Rev.* **D 56** (1997) 5867 [[hep-ph/9705258](#)].

- [18] A. Bialas and W. Szeremeta, *Heavy quark production by double pomeron exchange*, *Phys. Lett.* **B 296** (1992) 191;
A. Bialas and R.A. Janik, *Rapidity and transverse momentum distributions of heavy quarks produced by double pomeron exchange*, *Z. Physik* **C 62** (1994) 487.
- [19] J. Pumplin, *Two gluon exchange model predictions for double pomeron jet production*, *Phys. Rev.* **D 52** (1995) 1477 [[hep-ph/9412381](#)].
- [20] F. Yuan, *Heavy quarkonium production in double pomeron exchange processes in perturbative QCD*, *Phys. Lett.* **B 510** (2001) 155 [[hep-ph/0103213](#)].
- [21] S. Kim, *Decay rates of various bottomonium systems*, *Nucl. Phys.* **47** (Proc. Suppl.) (1996) 437.
- [22] PARTICLE DATA GROUP collaboration, D.E. Groom et al., *Review of particle physics*, *Eur. Phys. J.* **C 15** (2000) 1.
- [23] L. Randall and R. Sundrum, *A large mass hierarchy from a small extra dimension*, *Phys. Rev. Lett.* **83** (1999) 3370 [[hep-ph/9905221](#)].
- [24] M. Chaichian, A. Datta, K. Huitu and Z.-h. Yu, *Radion and Higgs mixing at the LHC*, *Phys. Lett.* **B 524** (2002) 161 [[hep-ph/0110035](#)].
- [25] M.G. Albrow and A. Rostovtsev, *Searching for the Higgs at hadron colliders using the missing mass method*, FERMILAB-PUB-00-173. Aug. 2000.